

## Chapter Three

### Reactions of Nitric Oxide with the Reduced Non-Heme Iron Center of the Soluble Methane Monooxygenase Hydroxylase

#### Introduction

The soluble methane monooxygenase (sMMO) enzyme system in methanotrophic bacteria catalyzes the oxidation of methane to methanol and water by using dioxygen as an oxidant at a non-heme carboxylate-bridged diiron center (1-5). One atom of the dioxygen molecule is incorporated into the product alcohol and the other is released as water. Two electrons from NADH and two protons are also required. The controlled oxidation of hydrocarbons by this and similar systems is a topic of considerable current interest.

The sMMO system has three known protein components. The hydroxylase (H) is a homodimer ( $\alpha_2\beta_2\gamma_2$ ) that contains a non-heme diiron center in each of its  $\alpha$  subunits. The iron center is the locus of dioxygen activation and methane hydroxylation. The reductase (R) is responsible for transferring electrons from NADH through protein-bound FAD and [2Fe-2S] centers to the diiron center in the hydroxylase. A third component, protein B or the coupling protein, serves to couple this electron transfer to alkane hydroxylation.

The catalytic cycle of the hydroxylase enzyme has been examined in detail by using a variety of rapid kinetics methodologies, including optical stopped-flow and freeze-quench Mössbauer spectroscopy. Typically, the iron center is chemically reduced to the diiron(II) state, designated  $H_{red}$ , and mixed rapidly with solutions containing dioxygen. The first well characterized intermediate after  $H_{red}$  is  $H_{peroxo}$  (6, 7), assigned as a ( $\mu$ -1,2-peroxo)diiron(III) species through

spectroscopic comparison with crystallographically characterized model compounds (8-10). The H<sub>peroxo</sub> intermediate spontaneously transforms into intermediate Q (7, 11), which has been assigned a di( $\mu$ -oxo)diiron(IV) structure based on EXAFS evidence (12), although other interpretations of the data are possible (1). Intermediate Q reacts with alkane substrates to form hydroxylated products, returning the catalytic center to its resting, diiron(III) state, designated H<sub>ox</sub>.

The role of protein B is not well understood. The catalytic cycle described above does not take place if protein B is absent. The diiron(II) center is instead oxidized to the diiron(III) state without observable accumulation of the H<sub>peroxo</sub> and Q intermediates. Under steady state conditions, protein B from the *M. capsulatus* (Bath) sMMO system has a dramatic effect on turnover (13, 14). When it is not present, hydroxylation of products is completely uncoupled from dioxygen and NADH consumption. Instead, dioxygen is reduced to water. As protein B is added to the system, the coupling of hydroxylated product formation to dioxygen consumption increases. At a 1:1 ratio of protein B to hydroxylase the system is fully coupled. As the ratio of protein B to hydroxylase increases beyond 1, the hydroxylation reaction is inhibited. These observations are consistent with protein B having a role as a regulatory protein. The coupling protein also has a direct effect on the iron center. The presence of protein B causes the g = 16 integer spin EPR signal of H<sub>red</sub> from *M. trichosporium* OB3b to become sharper and more intense (15). Magnetic circular dichroism experiments indicate that protein B effects a major coordination change at one of the iron atoms in the active site (16, 17), the details of which are obscure. Knowledge of these changes is crucial both structurally modeling and comprehending the mechanism of the sMMO system.

Nitric oxide is a useful surrogate for dioxygen (18) and reacts similarly with iron to afford complexes having useful spectroscopic properties. Species produced by the reaction of iron(II) with nitric oxide typically are brightly colored,  $S = 3/2$  systems (19-21) with characteristic EPR signals (22-24), and are designated as  $\{\text{Fe}(\text{NO})\}^7$  (21), where the superscript refers to the sum of d electrons from iron and  $\pi^*$  electrons from the NO molecule.

Nitric oxide has been useful as a probe of non-heme, carboxylate-bridged diiron centers. In hemerythrin (Hr), one NO molecule binds to a single iron atom (25, 26). The resulting  $\{\text{Fe}(\text{NO})\}^7$  center couples antiferromagnetically with the other high-spin iron(II) center to form an EPR-active,  $S = 1/2$  spin system. The mode of NO binding was proposed to be very similar to that of  $\text{O}_2$ , with the uncoordinated oxygen atom of the NO ligand forming a hydrogen bond with the hydroxide bridge present in deoxyhemerythrin. This proposal was supported by a shift in the Raman spectrum due to changes in the Fe–N–O bending mode upon deuteration of the bridge. In the R2 subunit of ribonucleotide reductase, a more complex, heterogeneous reaction was observed (27). Addition of NO to the diiron(II) center of this protein afforded a complex having optical features at 450 and 620 nm. Mössbauer spectra of the product revealed a mixture comprising 77% of a diiron dinitrosyl species,  $[\{\text{Fe}(\text{NO})\}^7]_2$ , 13% of mononuclear  $\{\text{Fe}(\text{NO})\}^7$ , and 10% of an unidentified species. The two  $S = 3/2$   $\{\text{Fe}(\text{NO})\}^7$  centers in the diiron dinitrosyl adduct were antiferromagnetically coupled to form an EPR-silent species, which decomposed with first order kinetics, reductively coupling the two NO molecules to form  $\text{N}_2\text{O}$  and oxidized RNR-R2. This diiron dinitrosyl species was proposed to be a model of the peroxy intermediate in the RNR-R2 reaction cycle. The spectroscopic properties of these and selected model compounds with carboxylate-bridged, non-heme diiron centers complexed with NO are listed in Table 1.

The present article describes a study of the reactions of NO with the reduced diiron center of the sMMO hydroxylase from *M. capsulatus* (Bath). These experiments afford the opportunity not only to model the O<sub>2</sub> binding step, but also to probe the role of protein B. The results provide important information about how the coupling protein can alter the dinuclear iron center for reaction with dioxygen.

## Experimental Procedures

*Preparation and Purification of sMMO Hydroxylase (H) and Protein B.* Samples of sMMO hydroxylase and of <sup>57</sup>Fe-enriched sMMO hydroxylase were prepared from *Methylococcus capsulatus* (Bath) as previously described (28). Protein B was prepared from a cloned source in *E. coli* as reported (29). Purity of the components was assured by SDS-PAGE electrophoresis. Specific activity and iron content were measured as described and were in the range of 250-350 nmol propylene formed mg<sup>-1</sup> min<sup>-1</sup> and 3.4-3.7 Fe/protein, respectively.

*Preparation of Diiron(II) sMMO Hydroxylase (H<sub>red</sub>).* Purified sMMOH samples were prepared in 25 mM MOPS pH 7.0 with stoichiometric amounts of the mediator methyl viologen. The sample was degassed by 15-20 cycles of vacuum evacuation followed by argon backfilling. A degassed solution of ~10 mM sodium dithionite in 50 mM MOPS (pH 8.6) buffer was prepared and used to titrate the protein solution until the methyl viologen changed from yellow to blue. A 30 min time period was sufficient for complete reduction. The sample was then anaerobically dialyzed against two successive 500 mL volumes of 25 mM MOPS pH 7.0 to remove the mediator and reductant.

*Addition of NO Gas.* H<sub>red</sub> samples were prepared and placed in vials sealed with rubber septa both with and without the presence of 2 equiv of protein B.

Nitric oxide (99+, Matheson) was purified by passage through a concentrated NaOH solution to remove NO<sub>2</sub>. The NO gas was passed over the protein solution for 5 min with manual agitation to ensure complete reaction. A change from colorless to green-brown was immediately observed. The NO stream was replaced by an argon stream for 10 min to remove excess NO. Samples for EPR and Mössbauer were anaerobically transferred to appropriate cells and frozen in liquid nitrogen approximately 45 min after NO addition.

*Addition of NO via NONOates.* NONOates are compounds that are stable at high pH, but decompose to NO gas at pH 7 (30, 31). They are useful as a means for delivering precisely known amounts of NO in aqueous solutions. NONOates are stored as concentrated stocks in 10 mM NaOH. The rate of decomposition was determined by introducing a small volume of concentrated NONOate into a 25 mM MOPS pH 7.0 solution and observing the disappearance of the NONOate UV absorption in mock reactions. This known rate of decomposition was used to determine how much NO is present in a protein solution at any given time after introduction of the NONOate.

H<sub>red</sub> samples were prepared and placed in septum-sealed vials both with and without the presence of 2 equiv of protein B. A small amount (< 1% of sample volume) of concentrated diethylamine-NONOate (Cayman Chemicals) was added, and the reaction was allowed to proceed until the desired amount of NO was generated. Samples were frozen in EPR tubes or Mössbauer cells.

*Stopped-flow Spectrophotometry.* Stopped-flow experiments were performed to observe the reaction of H<sub>red</sub> with nitric oxide both with and without the presence of 2 equiv of protein B. This ratio of B to H was chosen for direct comparison with previously reported stopped-flow studies with dioxygen (11). Protein samples, typically 50-150 μM in H<sub>red</sub>, were loaded anaerobically into Hamilton gastight syringes with Luer-lock fittings. Saturated NO solutions were

prepared by first saturating 25 mM MOPS pH 7.0 solutions with argon gas to remove O<sub>2</sub>. The solutions were cooled to 4 °C and NO gas was bubbled through them for 15-30 min, resulting in a solution of approximately 2.9 mM NO. This solution was diluted to the desired NO concentration with anaerobic buffer and transferred to a Hamilton gastight syringe. Both the protein- and gas-containing syringes were used to load the drive syringes of a HiTech SF61 DX2 stopped-flow instrument. Both syringes were allowed to equilibrate in a constant temperature bath for 15-30 min before recording data. Two detection systems were used. A HiTech UV-vis diode array spectrophotometer collected data over a wide range of wavelengths. A photomultiplier tube and monochrometer were used to collect data at single wavelengths.

To examine the effects of methane substrate, the NO containing buffer was further treated. An anaerobic methane stock solution was prepared by bubbling argon followed by methane through a septum sealed vial of 25 mM MOPS pH 7.0. An aliquot of this solution was mixed in a 1:1 ratio with NO-saturated buffer before being loaded into the stopped-flow syringe.

*Rapid Freeze-quench Mössbauer Sample Preparation.* All freeze-quench samples were made as described in detail elsewhere (32). Rapid freeze-quench samples were performed with an Update instruments Model 705A computer and syringe-ram apparatus. Protein solutions were 1.1 mM in H<sub>red</sub>. NO solutions were prepared by purging a 25 mM MOPS buffer sample with argon and allowing it to come to equilibrium in a chamber filled with 1 atm of NO gas at room temperature. The final NO concentration was approximately 1.9 mM. The reactants were loaded into RFQ syringes and allowed to cool to 4 °C in an ice/water bath. The protein and gas solutions were mixed rapidly, allowed to react for various fixed time periods governed by the length of tubing between the

mixing chamber and the nozzle, and then sprayed into isopentane at -140 °C. The resulting snow was packed into Delrin® Mössbauer sample holders.

*Rapid Freeze-quench EPR Sample Preparation.* Rapid freeze-quench samples were prepared by using an Update Instruments Model 715 syringe-ram apparatus. Solutions of proteins and NO gas were prepared in a manner similar to that described above for the stopped-flow procedures. H<sub>red</sub> concentrations were approximately 300 μM. The reactants were loaded into gas tight syringes and cooled to 4 °C. The solutions were mixed rapidly, allowed to react for fixed time periods (28 ms - 60 s) in the aging tubing, and then sprayed into isopentane at approximately -140 °C. The resulting snow was packed into quartz EPR tubes.

*EPR Measurements.* X-band EPR spectra were measured on a Bruker ESP 300 spectrometer equipped with an Oxford EPR 900 liquid helium cryostat. Spectra were recorded under the following conditions: temperature, 4.2 - 8.0 K; microwave frequency, 9.41 - 9.65 GHz; microwave power, 10 μW - 158 mW; modulation frequency, 100 kHz; modulation amplitude, 6.64 Gauss. Specific conditions are reported on the individual spectra. EPR quantitation was performed by double integration under nonsaturating conditions using 0.960 mM CuSO<sub>4</sub>, 1 M NaClO<sub>4</sub> as a standard. Quantitations were corrected for g-value anisotropy and spin. EPR power saturation data were collected by measuring signal intensity as a function of the applied microwave power over a range of four orders of magnitude.

*Mössbauer Measurements.* Mössbauer spectra were collected by using either a weak-field spectrometer equipped with a Janis 8DT variable-temperature cryostat or a strong-field spectrometer outfitted with a Janis 12 CNDT/SC SuperVaritemp cryostat encasing an 8 T superconducting magnet. Both spectrometers were operated in a constant acceleration mode in a transmission

geometry. The centroid of a room temperature iron foil spectrum was used as the zero velocity reference point.

*Nitrous Oxide Detection by Gas Chromatography.* Solutions of H<sub>red</sub> both with and without 2 equiv of protein B were prepared in 1 mL septum-sealed vials. These samples were ~300 μM in H<sub>red</sub> and 300 μL in volume. A saturated 2.9 mM NO buffer sample was prepared at 4 °C as described above. A 100 μL aliquot of the NO buffer was added to the protein sample. At 5 min and 30 min intervals after NO addition, 50 μL headspace aliquots were removed and injected on a Hewlett-Packard 5890 gas chromatograph equipped with a 6 ft. packed Haysep column and a thermal conductivity detection system. The column He gas flow rate was 21-23 mL/min at 35 °C. Under these conditions, retention times for various gases were determined to be ~1.3 min for N<sub>2</sub> and NO and 5.1-5.2 min for N<sub>2</sub>O. N<sub>2</sub>O standards were prepared by assembling mock reaction vials with BSA at w/v concentrations equivalent to the hydroxylase in 25 mM MOPS buffer. Known amounts of N<sub>2</sub>O were injected into these vials and they were subjected to the same 5 min and 30 min 50 μL headspace analysis as the H<sub>red</sub>+NO samples. Several N<sub>2</sub>O vials were analyzed and a standard curve was constructed.

## Results

*Preparation of NO complexes of H<sub>red</sub> and H<sub>red</sub>+2B.* Addition of NO gas to samples of H<sub>red</sub> and H<sub>red</sub> with two equiv of protein B (H<sub>red</sub>+2B) immediately afforded greenish brown solutions, henceforth designated H<sub>red</sub>+NO and H<sub>red</sub>+2B+NO, respectively. The NO addition method was varied and included direct addition of excess NO gas, direct addition of limited quantities of NO gas by a gas-tight Hamilton syringe, addition of NO via a NONOate precursor and addition of NO by NO-saturated buffer. No appreciable difference was observed

in the resulting optical or EPR spectral results for samples generated by these different methods.

*Optical Absorption Spectra.* The optical spectrum of the H<sub>red</sub>+NO adduct is shown in Figure 1, with maxima at 450 nm and 620 nm. The extinction coefficient at 450 nm was determined to be ~740 M<sup>-1</sup> cm<sup>-1</sup>. The spectrum of the H<sub>red</sub>+2B+NO complex was nearly identical, the only difference being a small decrease in  $\epsilon$  to approximately ~700 M<sup>-1</sup> cm<sup>-1</sup>.

These spectra are quite similar to those of the dinitrosyl adduct of [Fe<sub>2</sub>(μ-XDK)(μ-O<sub>2</sub>CPh)(ImH)<sub>2</sub>(O<sub>2</sub>CPh)(MeOH)] (33, 34), also shown in Figure 1. This compound has a ligand composition identical to that of H<sub>red</sub> and displays similar reactivity toward dioxygen. Colored intermediates with characteristic peroxy-to-iron charge transfer bands form rapidly upon reaction with dioxygen. This compound forms a dinitrosyl adduct upon reaction with NO, as established by Mössbauer and EXAFS spectroscopy (34).

*Mössbauer Analysis of H<sub>red</sub>+NO and H<sub>red</sub>+2B+NO Samples.* Figure 2 shows the zero-field Mössbauer spectra recorded at 4.2 K of samples of H<sub>red</sub> and H<sub>red</sub> with 2 equiv. of protein B that has been allowed to react with NO for approximately 45 min. Several species can be identified, the relative proportions of which are summarized in Table 2. Both unreacted H<sub>red</sub> and H<sub>ox</sub> can be identified in the spectra according to their previously identified characteristics (11, 28). Species consistent with mononuclear {Fe(NO)}<sup>7</sup> and mononuclear Fe(III) were identified on the basis of their isomer shift and high field Mössbauer properties. A diamagnetic species with Mössbauer parameters consistent with the dinitrosyl adduct of H<sub>red</sub>, termed H<sub>dinitrosyl</sub>, was detected. Finally, an unknown diamagnetic species with a very low isomer shift, 0.2 mm/s, was identified. This low isomer shift is consistent with an Fe(IV)-containing compound, such as species containing an Fe(NO)<sub>2</sub> unit.

*EPR Analysis of  $H_{red}+NO$  and  $H_{red}+2B+NO$  Samples.* Figure 3 shows the X-band EPR spectra of the  $H_{red}+NO$  and  $H_{red}+2B+NO$  adducts at 4-5 K. The major signal in the  $H_{red}+NO$  spectra has  $g = 4.08, 4.02$  and  $2.00$  and arises from an  $S = 3/2$  species consistent with mononuclear  $\{Fe(NO)\}^7$  as observed in the Mössbauer spectra described above. Determination of the spin concentration of this signal from several independent samples showed that it represents 20-30% of the total iron in the sample. The half-saturation value,  $P_{1/2}$ , for this  $\{Fe(NO)\}^7$  center was determined to be  $\sim 7$  mW. Interestingly, this signal splits into at least three species with slightly different rhombicities ( $g = 4.08, 4.02; g = 4.15, 3.95$ ; and  $g = 4.22, 3.86$ ) upon addition of protein B. This splitting was obtained by addition of protein B either before or after NO exposure. Mononuclear Fe(III) associated with exogenous iron impurities is present, also consistent with the Mössbauer samples, exhibiting a  $g = 4.3$  high spin ferric signal. Spin quantitation of this signal reveals that it represents 2-6% of total iron in the sample.

*Stopped-flow Analysis of the  $H_{red}+NO$  and  $H_{red}+2B+NO$  Reaction.* Stopped-flow optical spectrophotometry was used to analyze the kinetics of formation and decay of species formed in the reaction of NO with the reduced diiron center of the hydroxylase. As shown in Figure 4, an optical signal with maxima at 450 nm and 620 nm builds up rapidly upon mixing, maximizing at  $\sim 150$  ms. After 150 ms, the signal decays slowly until all visible changes stop, approximately 20-30 min after NO addition. At that time, the signal had approximately half of its 150 ms maximum intensity. The buildup was best fit by a biphasic  $A \rightarrow B \rightarrow C$  model that, at  $4$  °C, had rate constants  $k_1 = 78 \pm 6$  s $^{-1}$  and  $k_2 = 18 \pm 2.7$  s $^{-1}$ . Global analysis of the stopped-flow data revealed that the spectra of species B and C in the two phase model were nearly identical, the only difference being a 1-2% decrease in extinction coefficient for B. The decay of the optical signal could not be fit well to any specific model, indicating that it is a complex process

comprising various component reactions. First-order fits approximating the decay at 4 °C produced a rate constant of  $0.06 \pm 0.02 \text{ s}^{-1}$ . Variation of the NO concentration over a 10-fold range did not appreciably change the observed rate constants. All stopped-flow experiments subsequently reported here were conducted with at least a 20-fold excess of NO. The temperature dependence of the rate constants for buildup of the optical species was measured. Eyring plots of these data, shown in Figure 5, yielded the activation parameters shown in Table 3. A simulation of the time-dependent growth and decay of species formed in the  $\text{H}_{\text{red}} + \text{NO}$  reaction, obtained by using the rate constants obtained from stopped-flow data, is presented in Figure 6.

The reaction of NO with the reduced hydroxylase and two equiv of protein B was also examined by stopped-flow spectroscopy. The optical signals detected were essentially identical to those in the  $\text{H}_{\text{red}} + \text{NO}$  reactions, the major difference being a small diminution in observed extinction coefficient as described above. The buildup of this species could, in contrast with the  $\text{H}_{\text{red}} + \text{NO}$  reactions, be fit well by a single exponential buildup. At 4 °C, the rate constant was determined to be  $k = 26 \pm 0.8 \text{ s}^{-1}$ . The decay, as in the  $\text{H}_{\text{red}} + \text{NO}$  case, was too complex to be fit satisfactorily by a simple model. First order approximation of the decay yielded a rate of  $0.05 \pm 0.02 \text{ s}^{-1}$ , within error the same value as for the  $\text{H}_{\text{red}} + \text{NO}$  reaction. Temperature-dependent studies of this reaction produced activation parameters listed in Table 3, with the Eyring plots of these data shown in Figure 5.

The kinetics of the  $\text{H}_{\text{red}} + \text{NO}$  and  $\text{H}_{\text{red}} + 2\text{B} + \text{NO}$  reactions were also examined in the presence of methane. The optical bands that appeared were identical to those formed without methane present. The rates for both reactions could be fit well by first order equations.

*Rapid Freeze-Quench EPR Characterization of H<sub>red</sub>+NO Reaction Species.*

Freeze-quench samples of the H<sub>red</sub>+NO reaction were collected at 28 ms, 158 ms, 1 s and 60 s. Spectra taken from those samples are shown in Figure 7. The signal corresponding to the S = 3/2 {Fe(NO)}<sup>7</sup> species can be seen growing in at 28 ms, achieving its maximal value in the 158 ms sample. Fits of these data indicate that the first order rate constant for the buildup of this signal is 10-20 s<sup>-1</sup> at 4 °C. Its concentration, 27% of the total amount of iron centers in the sample, does not change within the error limits of the experiment after this time point (see inset, Figure 7). The signal at g = 1.97 is due to free NO in solution, and can be observed to decrease in intensity during the time course. In the 60 s sample, a signal at g = 2.04 has appeared. This signal is most likely due to a small amount of some S = 1/2, Fe(NO)<sub>2</sub> species. It accounts for <2% of total iron in the sample.

*Mössbauer Characterization of H<sub>red</sub>+NO Reaction Species.* The protein concentration used in these experiments was much higher than in other H<sub>red</sub>+NO work, in order to optimize the signal-to-noise ratio obtainable with Mössbauer spectroscopy. A consequence of this limitation is that the NO concentration (~1.9 mM) is stoichiometric with respect to the amount of iron (~3.9 mM). As a result, more than 50% of the hydroxylase remains in the reduced state. In addition, the rates of the NO reactions cannot be directly compared with those obtained in the stopped-flow and EPR freeze-quench experiments, because those data were obtained under pseudo-first-order conditions where NO was in great excess.

Figure 8 shows the Mossbauer spectra of the reaction of H<sub>red</sub> with NO at three different times after mixing, and Table 4 contains the Mössbauer parameters used for the fits to the data. Reactions were quenched at 61 ms, 126 ms, and 440 ms. The major component of all the spectra is a quadrupole doublet attributable to unreacted H<sub>red</sub>. A ~10% amount of H<sub>ox</sub> can be seen in all samples

that cannot be reduced, as previously observed for the *M. capsulatus* (Bath) hydroxylase (11). These two species can be identified in the spectra according to their previously identified characteristics (11, 28). A species corresponding to mononuclear  $\{\text{Fe}(\text{NO})\}^7$  develops very quickly, reaching its maximal concentration before 61 ms. It does not decay after 440 ms, in accord with the freeze-quench EPR data. A species with Mössbauer parameters characteristic of  $\text{H}_{\text{dinitrosyl}}$  also forms very quickly, consistent with the stopped-flow evidence. This species decays to approximately 80% of its maximal value at the 440 ms time point. This rate of decay is slightly faster than that observed in the stopped-flow experiments. Such a difference might arise if the “unidentified” decay product of  $\text{H}_{\text{dinitrosyl}}$  has optical properties similar to that of other iron nitrosyl species, a likely possibility. Thus the measured decay rate of  $\text{H}_{\text{dinitrosyl}}$  as observed by optical stopped-flow spectroscopy may be inaccurate owing to the buildup of other colored components. The Mössbauer parameters of a decay product from the  $\text{H}_{\text{red}} + \text{NO}$  reaction at the 440 ms time point suggest that it may be some  $\text{Fe}(\text{NO})_2$  containing species.

The Mössbauer freeze-quench work serves to identify  $\text{H}_{\text{dinitrosyl}}$  species as the diamagnetic reaction component that forms and decays slowly in the stopped-flow experiment (Figure 6). It also supports the rapid freeze-quench EPR data indicating that a mononuclear  $\{\text{Fe}(\text{NO})\}^7$  species develops very quickly upon reaction of NO with  $\text{H}_{\text{red}}$  and does not decay.

*Decay of  $\text{H}_{\text{dinitrosyl}}$  to  $\text{N}_2\text{O}$ .* The dinitrosyl adduct of the diiron center in the R2 subunit of ribonucleotide reductase decays slowly with a first order rate constant of  $\sim 0.013 \text{ min}^{-1}$  to  $\text{R2}_{\text{met}}$  and  $\text{N}_2\text{O}$  (27). To determine whether the sMMO dinitrosyl adduct undergoes similar chemistry, the headspace of a  $\text{H}_{\text{red}} + \text{NO}$  sample was analyzed for  $\text{N}_2\text{O}$  by gas chromatography. Samples taken at both 5 and 30 min after exposure to NO contained 0.06-0.12 equiv of  $\text{N}_2\text{O}$  per

hydroxylase diiron center. Similar samples taken of the  $H_{red} + 2B + NO$  reaction contained 0.10-0.16 equiv of  $N_2O$  per hydroxylase diiron center. One of the decay pathways of the  $H_{dinitrosyl}$  species is reductive elimination of  $N_2O$ , most likely affording  $H_{ox}$  as a product.

## Discussion

*Reaction of Reduced Hydroxylase with NO to Form a Diiron Dinitrosyl Species.* Nitric oxide reacts with the reduced diiron center in the sMMO hydroxylase to afford a diiron dinitrosyl species referred to as  $H_{dinitrosyl}$ . As indicated in Figure 9, this EPR-silent  $\{Fe(NO)\}^7_2$  unit accounts for a majority of the iron, the remainder being mononuclear (vide infra). The  $H_{dinitrosyl}$  intermediate models  $H_{peroxy}$  in the sMMO reaction cycle. Its two  $\{Fe(NO)\}^7$  centers antiferromagnetically couple to afford a diamagnetic ground state.

The spectroscopic properties of  $H_{dinitrosyl}$  were revealed by stopped-flow, RFQ EPR and RFQ Mössbauer experiments. When  $H_{red}$  is allowed to react with NO, an optical spectrum rapidly grows in, maximizes after approximately 150 ms, and subsequently decays. The optical spectrum closely resembles those of diiron dinitrosyl model complexes (34) and a similar dinitrosyl adduct of the ribonucleotide reductase R2 protein (27). The optical data are therefore consistent with formation of a diiron dinitrosyl unit, but alone are insufficient to prove that assignment.

Rapid freeze-quench Mössbauer data indicate that only two species containing iron-nitrosyl moieties form within the first 500 ms of reaction. The isomer shifts and quadrupole splitting parameters are consistent with the formation of  $H_{dinitrosyl}$  and a mononuclear  $\{Fe(NO)\}^7$  center. Rapid-freeze-quench EPR as well as Mössbauer spectroscopy establish that the mononuclear  $\{Fe(NO)\}^7$

unit forms in the first 100 ms after NO addition and does not decay even after 60 s of reaction. The RFQ Mössbauer data reveal that the species assigned as H<sub>dinitrosyl</sub> decays after 150 ms. Since the stopped-flow experiments indicate that the optical bands at 450 and 620 nm bands also decay after 150 ms of reaction, it can be concluded that H<sub>dinitrosyl</sub> contributes significantly to the observed optical signals.

Although it is possible that the mononuclear {Fe(NO)}<sup>7</sup> species also absorbs at 450 and 620 nm, its contribution is likely to be minimal. The EPR spectra reveal that most of the iron is diamagnetic. Moreover, both the H<sub>dinitrosyl</sub> and mononuclear {Fe(NO)}<sup>7</sup> species appear, from all available data, to build up during the first 200 ms of reaction. If the contribution from a mononuclear {Fe(NO)}<sup>7</sup> species were similar to that of H<sub>dinitrosyl</sub>, one would expect to be able to distinguish the two formation rates kinetically. Instead, the stopped flow data could be fit well to simple A → B (H<sub>red</sub>+2B+NO) and A → B → C (H<sub>red</sub>+NO) models (vide infra). Fits to A → B, A → C buildup kinetics could not account for the data. It is therefore likely that only one major optical species forms in the H<sub>red</sub>+NO reaction. Since H<sub>dinitrosyl</sub> is a significant contributor to the optical signals seen in the stopped-flow spectroscopic experiments (vide supra), we conclude that H<sub>dinitrosyl</sub> is the major component.

*A Mononuclear {Fe(NO)}<sup>7</sup> Minority Species in the H<sub>red</sub>+NO Reaction.*  
Approximately 15-30% of the protein active sites react with NO to afford a paramagnetic, S = 3/2 {Fe(NO)}<sup>7</sup> unit. This species probably forms in the active site of the hydroxylase, because its EPR characteristics are affected by the presence of protein B (vide infra). We considered the possibility that the {Fe(NO)}<sup>7</sup> EPR signal might arise by reaction of the active sites to eject one iron atom, leaving the second to react with NO to form a mononuclear species. Such a mechanism is unlikely, however, since the iron atoms are well coordinated by

protein ligands and not expected be extruded from the active site simply by addition of nitric oxide. Moreover, the iron content of the hydroxylase was determined both before and following NO treatment, and no metal was lost as a result of the reaction (data not shown). The sMMO hydroxylase as currently isolated from *M. capsulatus* (Bath) has 3.4-3.7 iron atoms per molecule, fewer than the expected number of 4 if all sites were fully occupied. The absence of one iron atom from 20% of the active sites would amount to a 10% deficiency of total iron, consistent with the measured amount of iron in the protein. We therefore attribute the  $\{\text{Fe}(\text{NO})\}^7$  signal to mononuclear resulting from iron-depleted enzyme. Heterogeneity in the sMMO hydroxylase as isolated has previously been noted in kinetics and spectroscopic experiments (11).

*Effects of Protein B on the Mononuclear  $\{\text{Fe}(\text{NO})\}^7$  Species.* The presence of protein B causes the  $S = 3/2$  EPR signal of the  $\{\text{Fe}(\text{NO})\}^7$  species to split into three signals arising from three species with slightly different rhombicities. This phenomenon can be explained by the following model. The species with the least rhombic character ( $g = 4.08, 4.02$ ) is assigned to hydroxylase having no bound protein B. When protein B binds to this enzyme, the active site environment is altered, and the  $\{\text{Fe}(\text{NO})\}^7$  unit assumes a slightly more rhombic coordination environment. This species is assigned to the  $g = 4.15, 3.95$  signal. When a second molecule of protein B is bound, the iron site rhombicity is shifted even further, resulting in the observation of yet a third signal. Although interesting, this information is of limited value, for it is impossible to assign specific structures to the EPR observed species, and it is unlikely that the hydroxylase sites housing the  $\{\text{Fe}(\text{NO})\}^7$  centers have the native conformation.

*Comparison Between the Reactions of the sMMO Hydroxylase with NO and  $\text{O}_2$ .* The reaction of reduced hydroxylase with dioxygen in the absence of protein B has not been investigated in detail. Under these conditions, the consumption of

electrons and dioxygen is completely uncoupled from substrate hydroxylation (14), although the hydroxylase does become oxidized. Intermediates H<sub>peroxo</sub> and Q do not accumulate. By contrast, the reaction of H<sub>red</sub> and two equiv of protein B with O<sub>2</sub> has been well characterized (7, 11). The first step is the formation of the H<sub>peroxo</sub> intermediate, which is most likely a ( $\mu$ -1,2-peroxo)diiron(III) species. Further reaction produces intermediate Q, which oxidizes substrate. There are clear differences between the reactions with and without the presence of protein B.

The reaction of H<sub>red</sub> with NO to form the H<sub>peroxo</sub> analogue H<sub>dinitrosyl</sub>, however, occurs whether or not protein B is present. There are two possible models to explain this difference. If the binding of protein B to the hydroxylase modulates the iron-iron distance, in the absence of this component the iron atoms may not have the correct separation to accommodate a peroxide-bridged structure. If dioxygen were to bind under such circumstances, it might be reduced to the superoxo level but not react further without being released from iron. The binding of protein B might bring the iron atoms to the proper distance to support 1,2-peroxo bridge formation. With NO as the reactant, the iron-iron distance should not be an issue, and a dinitrosyl species should form whether or not protein B is present. In the second model, protein B adjusts the relative orientations of the available coordination sites in H<sub>red</sub>. Without protein B present, the coordination sites may be positioned such that peroxide cannot bridge. Binding of protein B may cause a shift of the ligands such that the available coordination sites are oriented to facilitate formation of a peroxide bridge. Such shifting would not be required for NO binding to the two iron atoms. These two models are not mutually exclusive.

The available crystallographic data for reduced sMMO hydroxylase (35) are consistent with aspects of both models, as illustrated in Figure 10. The iron-

iron distance in the diiron(II) center is 3.28 Å, shorter than the 4.0 Å distance in a crystallographically characterized  $\mu$ -1,2-peroxodiiron(III) model complex having the same Mössbauer parameters as  $H_{\text{peroxo}}$  (8). Moreover, there are two sites in the reduced diiron center structure that are occupied by very weakly bound water molecules, and are therefore available for exogenous ligand binding. These sites are approximately 80°, or gauche, to one another when viewed down the iron-iron axis, as illustrated in Figure 10. Binding of protein B may shift the coordination environment so as to bring these open coordination sites into a more eclipsed conformation and possibly also to alter the iron-iron distance. In Figure 11, this model is elaborated by hypothesizing that Glu243 undergoes a carboxylate shift, bridging the two iron atoms in a  $\mu$ -1,2 fashion, as previously hypothesized (35). The resulting diiron environment is very much like that in the ribonucleotide reductase R2<sub>red</sub> structure (36), where the available coordination sites are adjacent to one another. Details of this model may be altered when information about the structure of the complex between the hydroxylase and protein becomes available.

*Buildup of  $H_{\text{dinitrosyl}}$ .* The formation of  $H_{\text{dinitrosyl}}$  from the reaction of  $H_{\text{red}}$  with NO in the absence of protein B and methane is clearly biphasic. There are several possible interpretations of this result. The first is that the initial intermediate formed is a mononitrosyl species, with an NO molecule bound to one of the iron atoms. Reaction with a second NO would lead to formation of  $H_{\text{dinitrosyl}}$ . This hypothesis is not consistent with the available data, however. Such a mononitrosyl species would be expected to have spectroscopic properties very similar to those of the hemerythrin nitrosyl adduct. The ferrous iron and resulting {Fe(NO)}<sup>7</sup> center would couple antiferromagnetically, forming an S = 1/2 paramagnetic species that would be clearly visible by EPR spectroscopy in rapid freeze quench samples. No such signal was apparent. Moreover, the optical

spectrum of the intermediate revealed by global analysis of the stopped flow data closely resembles that of  $\text{H}_{\text{dinitrosyl}}$ . A mononitrosyl species would be expected to have a much smaller extinction coefficient.

The preferred explanation is that the initial reaction of the reduced diiron center with NO affords a dinitrosyl intermediate, designated  $[\text{H}_{\text{dinitrosyl}}]^*$ , which spontaneously rearranges to the final,  $\text{H}_{\text{dinitrosyl}}$ , form. Since the spectra of both species are nearly identical, they most likely have similar structures. It may be that the  $[\text{H}_{\text{dinitrosyl}}]^*$  to  $\text{H}_{\text{dinitrosyl}}$  transition is the result of a shift in the relative positions of the NO ligands, such as that drawn in Figure 11. Another possibility is that the iron ligand environments are largely unchanged, but that formation of the initial  $[\text{H}_{\text{dinitrosyl}}]^*$  species triggers a conformational change in the hydroxylase, producing  $\text{H}_{\text{dinitrosyl}}$ , which can be resolved kinetically.

The absence of an mononitrosyl intermediate in the reaction of NO with the reduced hydroxylase suggests a mechanism for the dioxygen reaction. This observation means that there is a readily available coordination site on each iron in the diiron active site that can quickly react with NO, resulting in the formation of the diiron dinitrosyl species without formation of a diiron mononitrosyl intermediate. This implies that the dioxygen reaction will proceed similarly, with near-simultaneous reaction of both oxygen atoms with the two iron atoms, respectively. This rules out a hemerythrin-type mechanism, where dioxygen binds to just one iron and is reduced to peroxide without coordination to the second iron.

When the  $\text{H}_{\text{red}} + \text{NO}$  reaction is performed in the presence of protein B the kinetics of buildup become monophasic; no  $[\text{H}_{\text{dinitrosyl}}]^*$  intermediate is observed. This observation can be interpreted in terms of the ligand rearrangement model (Figure 11). If the  $[\text{H}_{\text{dinitrosyl}}]^*$  to  $\text{H}_{\text{dinitrosyl}}$  transition as observed in reactions without protein B is due to a shift in NO ligands from gauche to eclipsed

conformations, the shift to monophasic kinetics upon addition of protein B is a reasonable consequence of that model.

The pseudo-first-order rate constant for reaction of NO with  $H_{red}+2B$  to form  $H_{dinitrosyl}$  at 4 °C is ~26 s<sup>-1</sup>. This value is nearly identical to the rate constant for reaction of O<sub>2</sub> with  $H_{red}+2B$  to form  $H_{peroxo}$  at 4 °C, ~25 s<sup>-1</sup> (6). The rate constants for the initial reaction with NO and O<sub>2</sub> with  $H_{red}$  alone cannot be compared, since no value is available for the latter reaction. The rate constant for initial reaction with NO (~78 s<sup>-1</sup>) is much larger than with protein B present, however, which suggests that the corresponding reaction with dioxygen would also be much faster.

An investigation of the  $H_{red}+NO$  reaction without protein B, but in the presence of methane, revealed monophasic buildup kinetic behavior, similar to that as observed in the presence of protein B. This result suggests that protein B and methane gas might be able to modify the sMMO hydroxylase structure in a similar way. For example, it may be that both protein B and methane serve to extrude weakly bound water from the diiron center, clearing the way for dioxygen and nitric oxide exogenous ligand binding. One of the two steps observed kinetically in the  $H_{red}+NO$  stopped-flow experiments might therefore be loss of bound water. Addition of methane or protein B would facilitate this reaction, resulting in first-order kinetics. A similar effect is well known in the cytochromes P-450 (37), where substrate binding to the protein facilitates loss of a water ligand from the heme cofactor. This explanation is also consistent with the activation parameters determined for the reactions of  $H_{red}+NO$  and  $H_{red}+2B+NO$ , as reported in Table 3. For the reaction of NO with  $H_{red}$  in the presence of protein B, a ΔS<sup>‡</sup> value of -24.9 cal mol<sup>-1</sup> K<sup>-1</sup> was calculated. This result is consistent with binding of two NO molecules to the iron center in the transition state. The corresponding reaction of NO with  $H_{red}$  without protein B

has a much less negative  $\Delta S^\ddagger$  value of  $-1.1 \text{ cal mol}^{-1} \text{ K}^{-1}$ . This value must reflect a compensating entropic gain in the transition state during the  $\text{H}_{\text{dinitrosyl}}$  formation, such as loss of water. These results support the hypothesis that protein B serves to organize the diiron cluster for optimal reaction with dioxygen, as shown in Figure 11. The substrate methane appears to share aspects of this organizational capacity.

*Decay of  $\text{H}_{\text{dinitrosyl}}$ .* The decay of the dinitrosyl species is complex and stopped-flow data for this process could not be fit well to any single model. The final mixture of products, as observed by Mössbauer spectroscopy, is heterogeneous (data not shown). In one decay pathway, nitrous oxide is eliminated from  $\text{H}_{\text{dinitrosyl}}$  to afford  $\text{H}_{\text{ox}}$ . This reaction is supported by gas chromatographic detection of  $\text{N}_2\text{O}$  in the headspace above the reaction mixture. The ability of protein B to increase the amount of  $\text{N}_2\text{O}$  formed is consistent with the hypothesis that, in  $\text{H}_{\text{dinitrosyl}}$ , the two NO molecules are oriented in positions unfavorable for N–N bond formation. Addition of protein B could reposition the two NO coordinated molecules towards one another, as depicted in Figure 11, to promote ligand migration and subsequent N–N bond formation. Reductive coupling between two metal-bound NO molecules to form  $\text{N}_2\text{O}$  is well precedented in the literature. The dinitrosyl adduct of the ribonucleotide reductase R2 subunit reacts in this manner (*vide supra*) (27).  $\text{N}_2\text{O}$  formation has also been observed in dicopper(I) complexes (38).

A second decay pathway is suggested by the presence of the “unidentified” species in the Mössbauer spectra of  $\text{H}_{\text{red}} + \text{NO}$  reaction mixtures. A product with a low isomer shift appears following decay of  $\text{H}_{\text{dinitrosyl}}$  in the freeze-quench Mössbauer data. This product is diamagnetic, has Mössbauer properties consistent with a high-valent Fe(IV) species, and may contain the  $\text{Fe}(\text{NO})_2$  fragment. The data are consistent with an active site having two such

$\{\text{Fe}(\text{NO})_2\}$  centers, antiferromagnetically coupled. Formation of such a species might result from further reaction of NO with the diiron center, displacing water or protein side chain ligands.

*Comparison with the NO Reactions of the Reduced Ribonucleotide Reductase R2 Subunit and Hemerythrin.* There are many similarities between the reactions of the reduced sMMO hydroxylase and R2<sub>red</sub> with NO. The major products formed by each enzyme are  $[\{\text{Fe}(\text{NO})\}^7]_2$  dinitrosyl species that model aspects of peroxy intermediates in the two systems. These dinitrosyl intermediates are metastable. In the R2 system, decay is slow and apparently consists only of reductive coupling to afford R2<sub>met</sub> and N<sub>2</sub>O. The H<sub>dinitrosyl</sub> adduct decays more rapidly, forming a more complex mixture of products. A reductive coupling pathway to form N<sub>2</sub>O is apparently present, but does not account for all of the H<sub>dinitrosyl</sub> decay.

The minor products formed in both reactions are also similar and include mononuclear  $\{\text{Fe}(\text{NO})\}^7$  centers and unidentified diamagnetic species having Mössbauer parameters consistent with  $[\{\text{Fe}(\text{NO})_2\}_2]$ . This species appears to be a byproduct of H<sub>dinitrosyl</sub> decay in the sMMO system. The similarities emphasize the close relationship between the structures and functions of these two enzymes, which is to perform chemistry by using dioxygen as a reactant. The NO chemistry contrasts with that of hemerythrin, which forms a relatively homogenous and stable NO adduct. The function of hemerythrin is to bind dioxygen reversibly, which is consistent with its reaction with NO.

The present results have been useful in suggesting specific roles for the action of protein B in the sMMO system. Examination of the NO chemistry of related proteins, including stearoyl Δ-9 desaturase (39, 40), toluene-4-monooxygenase (41) and phenol hydroxylase (42), would be useful in

elucidating the similarities and differences in the reactivities of this class of proteins.

## References

1. Valentine, A. M., and Lippard, S. J. (1997) *J. Chem. Soc., Dalton Trans.* 21, 3925-3931.
2. Kurtz, D. M. (1997) *JBIC* 2, 159-167.
3. Wallar, B. J., and Lipscomb, J. D. (1996) *Chem. Rev.* 96, 2625-2657.
4. Liu, K. E., and Lippard, S. J. (1995) in *Adv. Inorg. Chem.* (Sykes, A. G., Ed.) pp 263-289, Academic Press, San Diego.
5. Lipscomb, J. D. (1994) *Annu. Rev. Microbiol.* 48, 371-399.
6. Liu, K. E., Valentine, A. M., Qiu, D., Edmondson, D. E., Appelman, E. H., Spiro, T. G., and Lippard, S. J. (1995) *J. Am. Chem. Soc.* 117, 4997-4998.
7. Lee, S.-K., Nesheim, J. C., and Lipscomb, J. D. (1993) *J. Biol. Chem.* 268, 21569-21577.
8. Kim, K., and Lippard, S. J. (1996) *J. Am. Chem. Soc.* 118, 4914-4915.
9. Dong, Y., Yan, S., Young, V. G., and Que, L. (1996) *Agnew. Chem. Int. Ed. Engl.* 35, 618-620.
10. Ookubo, T., Sugimoto, H., Nagayama, T., Masuda, H., Sato, T., Tanaka, K., Maeda, Y., Okawa, H., Hayashi, Y., Uehara, A., and Suzuki, M. (1996) *J. Am. Chem. Soc.* 118, 701-702.
11. Liu, K. E., Valentine, A. M., Wang, D., Huynh, B. H., Edmondson, D. E., Salifoglou, A., and Lippard, S. J. (1995) *J. Am. Chem. Soc.* 117, 10174-10185.
12. Shu, L., Nesheim, J. C., Kauffmann, K., Münck, E., Lipscomb, J. D., and Que, L. (1997) *Science* 275, 515-517.
13. Green, J., and Dalton, H. (1985) *J. Biol. Chem.* 260, 15795-15801.
14. Gassner, G., and Lippard, S. J. (1998) *manuscript in preparation.*
15. Hendrich, M. P., Münck, E., Fox, B. G., and Lipscomb, J. D. (1990) *J. Am. Chem. Soc.* 112, 5861-5865.

16. Pulver, S., Froland, W. A., Fox, B. G., Lipscomb, J. D., and Solomon, E. I. (1993) *J. Am. Chem. Soc.* 115, 12409-12422.
17. Pulver, S. C., Froland, W. A., Lipscomb, J. D., and Solomon, E. I. (1997) *J. Am. Chem. Soc.* 119, 387-395.
18. Richter-Addo, G. B., and Legzdins, P. (1992) *Metal Nitrosyls*, Oxford Univeristy Press, New York.
19. Zhang, Y., Pavlosky, M. A., Brown, C. A., Westre, T. E., Hedman, B., Hodgson, K. O., and Solomon, E. I. (1992) *J. Am. Chem. Soc.* 114, 9189-9191.
20. Brown, C. A., Pavlosky, M. A., Westre, T. E., Zhang, Y., Hedman, B., Hodgson, K. O., and Solomon, E. I. (1995) *J. Am. Chem. Soc.* 117, 715-732.
21. Enemark, J. H., and Feltham, R. D. (1974) *Coord. Chem. Rev.* 13, 339-406.
22. Nelson, M. J. (1987) *J. Biol. Chem.* 262, 12137-12142.
23. Arciero, D. M., Lipscomb, J. D., Huynh, B. H., Kent, T. A., and Münck, E. (1983) *J. Biol. Chem.* 258, 14981-14991.
24. Arciero, D. M., Orville, A. M., and Lipscomb, J. D. (1985) *J. Biol. Chem.* 260, 14035-14044.
25. Nocek, J. M., Kurtz, D. M., Sage, J. T., Debrunner, P. G., Maroney, M. J., and Que, L. (1985) *J. Am. Chem. Soc.* 107, 3382-3384.
26. Nocek, J. M., Donald M. Kurtz, J., Sage, J. T., Xia, Y.-M., Debrunner, P., Shiemke, A. K., Sanders-Loehr, J., and Loehr, T. M. (1988) *Biochemistry* 27, 1014-1024.
27. Haskin, C. J., Ravi, N., Lynch, J. B., Münck, E., and Lawrence Que, J. (1995) *Biochemistry* 34, 11090-11098.
28. DeWitt, J. G., Bentsen, J. G., Rosenzweig, A. C., Hedman, B., Green, J., Pilkington, S., Papaefthymiou, G. C., Dalton, H., Hodgson, K. O., and Lippard, S. J. (1991) *J. Am. Chem. Soc.* 113, 9219-9235.

29. Liu, K. E., Johnson, C. C., Newcomb, M., and Lippard, S. J. (1993) *J. Am. Chem. Soc.* 115, 939-947.
30. Maragos, C. M., Morley, D., Wink, D. A., Dunams, T. M., Saavedra, J. E., Hoffman, A., Bove, A. A., Issac, L., Hrabie, J. A., and Keefer, L. K. (1991) *J. Med. Chem* 34, 3242-3247.
31. Morley, D., and Keefer, L. K. (1993) *J. of Card. Pharm.* 22, S3-S9.
32. Ravi, N., Bollinger, J. M., Huynh, B. H., Edmundson, D., and Stubbe, J. (1994) *J. Am. Chem. Soc.* 116, 8007-8014.
33. Herold, S., and Lippard, S. J. (1997) *J. Am. Chem. Soc.* 119, 145-156.
34. Mizoguchi, T. J., DuBois, J. L., Bautista, M. T., Hedman, B., Hodgson, K. O., and Lippard, S. J. (1998) *manuscript in preparation*.
35. Rosenzweig, A. C., Nordlund, P., Takahara, P. M., Frederick, C. A., and Lippard, S. J. (1995) *Chemistry & Biology* 2, 409-418.
36. Åberg, A. (1993), Stockholm University.
37. Sono, M., Roach, M. P., Coulter, E. D., and Dawson, J. H. (1996) *Chem. Rev.* 96.
38. Paul, P. P., and Karlin, K. D. (1991) *J. Am. Chem. Soc.* 113, 6331-6332.
39. Fox, B. G., Shanklin, J., Somerville, C., and Münck, E. (1993) *Proc. Natl. Acad. Sci. USA* 90, 2486-2490.
40. Fox, B. G., Shanklin, J., Ai, J., Loehr, T. M., and Sanders-Loehr, J. (1994) *Biochemistry* 33, 12776-12786.
41. Pikus, J. D., Studts, J. M., Achim, C., Kauffmann, K. E., Münck, E., Steffan, R. J., McClay, K., and Fox, B. G. (1996) *Biochemistry* 35, 9106-9119.
42. Nordlund, I., Powłowski, J., and Shingler, V. (1990) *J. Bact.* 172, 6826-6833.
43. Feig, A. L., Bautista, M. T., and Lippard, S. J. (1996) *Inorg. Chem.* 35, 6892-6898.
44. Kraulis, P. J. (1991) *J. Appl. Cryst.* 24, 946-950.

Table 1: Spectroscopic Properties of Non-Heme Carboxylate-Bridged Diiron Dinitrosyl Complexes.

Table 2:

Table 3: Activation Parameters from Eyring Plots of the Reaction of the Reduced Hydroxylase with Nitric Oxide.

Table 4: Time-Dependant Changes in the Mössbauer Spectra of Rapid Freeze-quench Time Points of the Reactions of Nitric Oxide with the Reduced sMMO Hydroxylase.

Figure 1: Optical spectra of the nitrosyl adduct of the reduced sMMO hydroxylase in 25 mM MOPS pH 7.0 (solid line) and the dinitrosyl adduct of  $[\text{Fe}_2(\mu\text{-XDK})(\mu\text{-O}_2\text{CPh})(\text{ImH})_2(\text{O}_2\text{CPh})(\text{MeOH})]$  in THF (dashed line) (34).



Figure 2: Mössbauer spectra of  $^{57}\text{Fe}$ -enriched  $\text{H}_{\text{red}} + \text{NO}$  and  $\text{H}_{\text{red}} + 2\text{B} + \text{NO}$ .  
Mössbauer spectra of  $^{57}\text{Fe}$ -enriched  $\text{H}_{\text{red}}$  (a) and  $\text{H}_{\text{red}}$  in the presence of two equiv. of protein B (b) reacted with NO for 45 min at 4 K in zero applied field. The solid line drawn through the data represent a simulation of the species detailed in Table 2.



Figure 3: EPR spectra of H<sub>red</sub> (a, b) and H<sub>red</sub> in the presence of two equiv of protein B (c) exposed to NO for 45 min. Spectra were obtained at 4.3-4.8 K, 100  $\mu$ W and 9.41 GHz. The buffer was 25 mM MOPS, pH 7.0.



Figure 4: (a) Buildup of the optical species associated with reaction of NO with  $H_{red}$  as observed by using stopped-flow spectroscopy at 4 °C. (b) Two comparative fits of the stopped-flow data. The first graph is a fit of the data to an  $A \rightarrow B$  buildup model and the second is a fit to an  $A \rightarrow B \rightarrow C$  buildup.



Figure 5: Eyring plots of the fast and slow reactions of the  $H_{red} + NO$   $H_{dinitrosyl}$  buildup and of the  $H_{red} + 2B + NO$   $H$  dinitrosyl buildup.



Figure 6: Simulation of the  $\text{H}_{\text{dinitrosyl}}$  formation and decay reactions by using rate constants derived from stopped-flow data. (top) Simulation of the first 500 ms of reaction. Also plotted is the total of the two dinitrosyl species, which corresponds to a fit of the stopped-flow optical data. (bottom) Simulation of the reaction from 100 ms to 10 s.



Figure 7: Time-dependent RFQ EPR spectra following the reaction of NO with the reduced sMMO hydroxylase at 4 °C. Spectra correspond to reaction times of 28 ms, 158 ms, 1 s and 30 s. Spectra were obtained at 4.5 K, 100 µW and 9.41 GHz. The inset is a plot of the concentration of the mononuclear  $\{\text{Fe}(\text{NO})\}^7$  species as a function of time, with the solid line showing a monophasic fit of the data.



Figure 8: Time-dependent Mössbauer spectra following the reaction of H<sub>red</sub> with NO at 4 °C. The H<sub>red</sub> Mössbauer spectrum has been subtracted from all time points. The solid line through the data corresponds to simulations using parameters listed in Table 4.



Figure 9: Proposed mechanism for the reactions of NO with H<sub>red</sub>.



Figure 10: Active site structure of the reduced sMMO hydroxylase. a. Ligand arrangement around the iron site. Note that the open coordination sites on each iron, occupied by weakly bound water ligands. b. Newman projection viewed down the Fe-Fe axis in the  $H_{red}$  active site. The positions of the open coordination sites are shown. These figures were prepared with MolScript (44).



Figure 11: Proposal for ligand movements in the active site of the reduced sMMO hydroxylase induced by binding of protein B. The schematic for the  $H_{red}$  active site is taken from the crystal structure of the reduced sMMOH, and the  $H_{red}+2B$  active site from the crystal structure of the reduced R2 subunit.

